

DAFTAR PUSTAKA

- Agami M, Waisel Y (1988) The role of fish in distribution and germination of seed of the submerged Macrophytes *Najas marina* L. and *Ruppia maritima* L. *Oecologia* 76, 83– 88.
- Ariyanti ES (2016) Dampak Perubahan Ekosistem Hutan Menjadi Agroforestri Karet, Kebun Karet, dan Kebun Kelapa Sawit Terhadap Keanekaragaman Jenis dan Kelimpahan Relatif Kelelawar : Studi Di Hutan Harapan PT.Restorasi Ekosistem Indonesia (REKI) dan Taman Nasional Bukit Duabelas, Jambi [tesis]. Bandar Lampung: Program Studi Ilmu Lingkungan. UniversitasLampung.
- Barnea A., Yomtov Y., Friedman J (1990) Differential germination of two closely related species of *Solanum* inresponse to bird ingestion. *Oikos* 57, 222–228.
- Bocchese RA., Oliveira AKM., Vicente EC (2007) Taxa e velocidade de germinação de sementes de *Cecropia pachystachya* Trécul (Cecropiaceae) ingeridas por *Artibeus lituratus* (Olfers, 1818) (Chiroptera: Phyllostomidae). *Acta Scientiarum Biological Science, Maringá*, v. 29, n. 4, p. 395-399, 2007.
- Bronstein JL, Hoffmann K (1987) Spatial and temporal variation in frugivory at a neotropical fig, *Ficus pertusa*. *Oikos* 49, 261-268.
- Bumrungsri S., Lang D., Harrower C., Sripaoraya E., Kitpith K., Racey PA (2013)The dawn bat, *Eonycteris spelaea* Dobson (Chiroptera: Pteropodidae) feeds mainly on pollen of economically important food plants in Thailand. *Acta Chiropterologica*, 15(1), 95-104.
- Chaveser IN., Rothenwohrer C., Tschapka M (2010) Dynamic feeding habits:efficiency of frugivory in a nectarivorous bat. *Can. J. Zool.* 88: 764-773
- Chaves- R HE., Urbano-N EA., Martinez-M DM., Zure D., Arboleda-V FA., Meluk-M H (2020) Endemic bats (Mamalia: Chiroptera) of Colombia: State of knowledge, distribution, and conservation. *Univ. Sci.* 25 (1): 55-94, 2020.
- Cl J., Fletcher C., Salim HM., Rahman A., Harrison RD., Potts MD (2011)Insectivorous bat assemblage in the hill dipterocarp forest of Temengor ForestReserve, Peninsular Malaysia. *Malayan Nature Journal*, 63(3), 569-576.
- Corbett GB, Hill JE (1980) A world list of mammalian species, British Museum ofNatural History. *Comstock Publishing, London*, 128, 1950-1959.
- Corlett RT (1990) Seed dispersal by the Lesser Short-nosed Bat (*Cynopterus brachyotis*, Pteropodidae, Megachiroptera). *Malayan Nature Journal*, 42:251-256.

- Dumont ER (2003) Bats and fruit: an ecomorphological approach. Pp. 389-429, *in*Bat ecology (T. H. Kunz and M.B. Fenton, eds.). The University of Chicago Press, Chicago, 779 pp.
- Findley JS (1993) Bats a Community Perspective Cambridge University Press, New York.
- Fleming TH, Williams CF (1990) Phenology, seed dispersal, and recruitment in *Cecropia peltata* (Moraceae) in Costa Rican tropical dry forest. *Journal of Tropical Ecology* 6, 163–178.
- Galindo GJ., Vazquez DG., Saldaña-V RA., Hernández-M JR (2009) A more efficient technique to collect seeds dispersed by bats. *Journal of Tropical Ecology*, 25(2), 205-209.
- Handley CO., Jr Wilson DE., Gardner AL (1991) Demography and natural history of the common fruit bat, *Artibeus jamaicensis*, on Barro Colorado Island, Panama. *Smithsonian Contributions to Zoology* 511, 1–173.
- Heer K., Albrecht L., Kalko EK (2010) Effects of ingestion by neotropical bats on germination parameters of native free-standing and strangler figs (*Ficus* sp., Moraceae). *Oecologia*, 163(2), 425-435.
- Heithaus ER., Fleming TH., Opler PA (1975) Foraging patterns and resource utilization in seven species of bats in a seasonal tropical forest. *Ecology*, 56(4), 841-854.
- Herbst LH (1986) The role of nitrogen fruit pulp in the nutrition of the frugivorous bat *Carollia perspicillata*. *Biotropica* 18, 39-44.
- Hodgkison R (2001) The ecology of fruit bats (Chiroptera: Pteropodidae) in a Malaysian lowland dipterocarp forest, with particular reference to the spotted-winged fruit bat (*Balionycteris maculata*, Thomas). Ph. D. dissertation, University of Aberdeen, Aberdeen, Scotland.
- Hodgkison R., Ayasse M., Kalko EKV., Haberlein C., Schulz S., Mustapha WAW., Zubaid A, Kunz TH (2007) Chemical ecology of fruit bat foraging behaviour in relation to the fruit odours of two species of paleotropical bat-dispersed figs (*Ficus hispida* and *Ficus scortechnii*). *J.Chem. Ecol.* 33: 2097-2110.
- Jim CY (2013) Sustainable urban greening strategies for compact cities in developing and developed economies. *Urban Ecosystems*, 16: 741–761.
- Kasso M, Balakrishnan M (2013) Ecological and economic importance of bats (Order Chiroptera). *ISRN Biodiversity*, 2013.
- Kingston T., Lim BL., Akbar Z (2006) *Bats of Krau wildlife reserve*. Penerbit Universiti Kebangsaan Malaysia.

- Kunz TH, S Parson (2009) Ecological Behavioral Methods For the Study of Bats. The Johns Hopkins University Press. Baltimore, United Stated of America.
- Lekagul, McNeely (1977) *Mammals of Thailand, Kurusapha*. Ladprao Press. Bangkok.
- Lim BL (1970) Food Habits and Breeding Cycle of the Malaysian Fruit-eating Bat, *Cynopterus brachyotis*. *Journal of Mammalogy* 51:174-177.
- Lisci M, Pacini E (1994) Germination ecology of drupelets of the fig (*Ficus carica* L.). *Botanical Journal of the Linnean Society* 114, 133–146.
- Lopez JE, Vaughan C (2004) Observations on the role of frugivorous bats as seed dispersers in Costa Rican secondary humid forests. *Acta Chiropterologica*, 6(1), 111-119.
- Lubis SH., Arifin HS., Samsoedin I (2013) Analisis cadangan karbon pohon pada lanskap hutan kota di DKI Jakarta. *Jurnal Penelitian Sosial dan Ekonomi Kehutanan*. 10: 1–20.
- Mayer AM, Poljakoff-Mayber A (1989) *The Germination of Seeds*, 4th edn. Pergamon Press, Oxford.
- Pradana DH, Tsang SM (2017) The role of fruit bats in plant community changes in an urban forest in Indonesia. *Raffles Bulletin of Zoology*, 65, 497-505.
- Prasetyo PN., Noerfahmy S., Tata HL (2011) Jenis-Jenis Kelelawar Khas Agroforest Sumatera. Bogor, Indonesia. World Agroforestry Center – ICRAF, SEA Regional Office. 75p.
- Ripperger S., Heymann WE., Tschapka M., Kalko VKE (2014) Fruit Characteristics Associated with Fruit Preferences in Frugivorous Bats and Saddleback Tamarins in Peru. *ECOTROPICA* 20: 53-64.
- Safi K., Meiri S., Jones KE (2013) Evolution of body size in bats. In: Smith FA & Lyons SK (eds.) *Animal Body Size: Linking Pattern and Process Across Space, Time, and Taxonomic Group*. University of Chicago Press, Chicago, USA. Pp. 95–115.
- Seidler TG, Plotkin JB (2006) Seed dispersal and spatial pattern in tropical trees. *PLoS Biology*, 4: 2132–2137.
- Soegiharto S., Kartono AP., Maryanto I (2001) The grouping of fruit bats based on pollen type characterized as food resource in Bogor Botanical Garden, Indonesia. *Jurnal Biologi Indonesia (Journal of Indonesian Biology)*, 6: 225-235.
- Suyanto A (2001) Seri Panduan Lapangan: Kelelawar di Indonesia. *Puslitbang- LIPI- Bogor*.
- Suyanto A (2002) Mamalia di Taman Nasional Gunung Halimun Jawa Barat. LIPI,JICA

and PHKA Bogor.

- Tan KH., Zubaid A., Kunz TH (1998) Food habits of Cynopterus brachyotis (Muller) (Chiroptera: Pteropodidae) in Peninsular Malaysia. *Journal of Tropical Ecology*, 14(3), 299-307.
- Tidemann CR (1987) Notes on the flying fox, Pteropus melanotus (Chiroptera:Pteropodidae) on Christmas Island, Indian Ocean. Australian Mammalogy 10, 89-91.
- Van Sommeren GRC (1972) Some fruit bat eat leaves. *Bull. East Afr. nat. Hist. Soc.*, February 1972:24-5.
- Voughan., Terry A., MR James., J Nicholas (2000) Mammalogy Fouth Edition. Harcourt College Publisher. United States of America. ISBN: 0-03-025034- X.
- Walker EP (1964) *Mammals of the world*, 2 vols. Baltimore: John Hopkins Press
- Yagihashi T., Hayashida M., Miyamoto T (1998) Effects of bird ingestion on seed germination of Sorbus commixta. *Oecologia* 114, 209–212.