

DAFTAR PUSTAKA

- Apriyani, N., Fanani, Z., & Almunady, A. (2019). *Uji Kualitas Sabun Padat Transparan dari Minyak Kelapa dan Minyak Kelapa Sawit Dengan Antikoksidan Ekstrak Likopen Buah Tomat*. Sriwijaya University.
- Atun, S. (2014). Metode Isolasi dan Identifikasi Struktur Senyawa Organik Bahan Alam. *Jurnal Konservasi Cagar Budaya Borobudur*, 8(2), 53–61.
- Baba, S., Chan, H. T., Kezuka, M., Inoue, T., & Chan, E. W. C. (2016). Artocarpus altilis and Pandanus tectorius: Two important fruits of Oceania with medicinal values. *Emirates Journal of Food and Agriculture*, 28(8), 531–539. <https://doi.org/10.9755/ejfa.2016-02-207>
- Boonphong, S., Baramee, A., & Kittakoop, P. (2007). Antitubercular and Antiplasmodial Prenylated Flavones from the Roots of Artocarpus altilis. *Chiang Mai Journal of Science*, 34(3), 339–344.
- Boonyaketgoson, S., Rukachaisirikul, V., & Phongpaichit, S. (2019). Deoxybenzoin and flavan derivatives from the twigs of Artocarpus lakoocha. *Phytochemistry Letters*, 31(March), 96–100. <https://doi.org/10.1016/j.phytol.2019.03.020>
- Boulebd, H. (2020). Comparative study of the radical scavenging behavior of ascorbic acid, BHT, BHA and Trolox: Experimental and theoretical study. *Journal of Molecular Structure*, 1201, 127210. <https://doi.org/10.1016/j.molstruc.2019.127210>
- Chen, C. C., Huang, Y. L., Ou, J. C., Lin, C. F., & Pan, T. M. (1993). Three new prenylflavones from Artocarpus altilis. *Journal of Natural Products*, 56(9), 1594–1597. <https://doi.org/10.1021/np50099a021>
- Chen, X., Liang, L., & Han, C. (2020). Borate suppresses the scavenging activity of gallic acid and plant polyphenol extracts on DPPH radical: A potential interference to DPPH assay. *Lwt*, 131, 109769. <https://doi.org/10.1016/j.lwt.2020.109769>
- Fang, S.-C., Hsu, C.-L., Yu, Y.-S., & Yen, G.-C. (2008). Cytotoxic Effects of New Geranyl Chalcone Derivatives Isolated from the Leaves of Artocarpus communis in SW 872 Human Liposarcoma Cells. 8859–8868.
- Guo, Y. Q., Tang, G. H., Lou, L. L., Li, W., Zhang, B., Liu, B., & Yin, S. (2018). Prenylated flavonoids as potent phosphodiesterase-4 inhibitors from Morus alba: Isolation, modification, and structure-activity relationship study. *European Journal of Medicinal Chemistry*, 144, 758–766. <https://doi.org/10.1016/j.ejmech.2017.12.057>
- Hafid, A. F., Aoki-Utsubo, C., Permanasari, A. A., Adianti, M., Tumewu, L., Widyawaruyanti, A., Wahyuningsih, S. P. A., Wahyuni, T. S., Lusida, M. I., Soetjipto, & Hotta, H. (2017). Antiviral activity of the dichloromethane extracts from Artocarpus heterophyllus leaves against hepatitis C virus. *Asian Pacific Journal of Tropical Biomedicine*, 7(7), 633–639. <https://doi.org/10.1016/j.apjtb.2017.06.003>
- Hakim, A., Jufri, A. W., Jamaluddin, Supriadi, & Mutmainnah, P. A. (2020). Understanding the Uniqueness of Artocarpus Flavonoids: Isolation and Structure Elucidation of Cycloartocarpin from the Roots of Artocarpus altilis. *Journal of Chemical Education*, 97(11), 4133–4136. <https://doi.org/10.1021/acs.jchemed.0c00199>

- Hennek, J., Alves, J., Yao, E., Goueli, S. A., & Zegzouti, H. (2016). *Bioluminescent kinase strips : A novel approach to targeted and flexible kinase inhibitor profiling*. 495. <https://doi.org/10.1016/j.ab.2015.11.007>
- Huang, Q., Zhong, Y., Li, J., Ye, Y., Wu, W., Chen, L., Feng, M., Yang, J., & Liu, S. (2020). Kinase inhibitor roscovitine as a PB2 cap-binding inhibitor against influenza a virus replication. *Biochemical and Biophysical Research Communications*, 526(4), 1143–1149. <https://doi.org/10.1016/j.bbrc.2020.04.034>
- Ishibashi, M. (2014). *Prenylated Flavonoids and Resveratrol Derivatives Isolated from Artocarpus communis with the Ability to Overcome TRAIL Resistance*. <https://doi.org/10.1021/np500734t>
- Jagtap, U. B., & Bapat, V. A. (2010). Artocarpus: A review of its traditional uses, phytochemistry and pharmacology. *Journal of Ethnopharmacology*, 129(2), 142–166.
- Kim, J. Y., Lee, W. S., Kim, Y. S., Curtis-long, M. J., Lee, B. W., Ryu, Y. B., & Park, K. H. (2011). Isolation of Cholinesterase-Inhibiting Flavonoids from Morus lhou. *Journal of Agricultural and Food Chemistry*, 59, 4589–4596.
- Kueté, V., Ango, P. Y., Fotso, G. W., Kapche, G. D. W. F., Dzoyem, J. P., & Wouking, A. G. (2011). *Antimicrobial activities of the methanol extract and compounds from Artocarpus communis (Moraceae)*. 2–6. <https://doi.org/10.1186/1472-6882-11-42>
- Kurniadewi, F., Dianhar, H., Mutiah, S., & Al-Liatsi, Z. Z. (2021). Two flavonoid derivatives from the wood of Nangkadak (Moraceae). *AIP Conference Proceedings*, 2331(April), 5–11. <https://doi.org/10.1063/5.0041840>
- Lan, W. C., Tzeng, C. W., Lin, C. C., Yen, F. L., & Ko, H. H. (2013). Prenylated flavonoids from Artocarpus altilis: Antioxidant activities and inhibitory effects on melanin production. *Phytochemistry*, 89, 78–88. <https://doi.org/10.1016/j.phytochem.2013.01.011>
- Lin, K., Liu, C., Tu, H., Ko, H., & Wei, B. (2009). Antioxidant prenylflavonoids from Artocarpus communis and Artocarpus elasticus. *Food Chemistry*, 115(2), 558–562. <https://doi.org/10.1016/j.foodchem.2008.12.059>
- Liu, Y., Yu, X., Zhang, W., Wang, T., Jiang, B., Tang, H., Su, Q., & Fu, Y. (2020). Prenylated chromones and flavonoids from Artocarpus heterophyllus with their potential antiproliferative and anti-inflammatory activities. *Bioorganic Chemistry*, 104030. <https://doi.org/10.1016/j.bioorg.2020.104030>
- Liyanaarachchi, G. D., Samarakkera, J. K. R. R., Mahanama, K. R. R., & Hemalal, K. D. P. (2018). Tyrosinase, elastase, hyaluronidase, inhibitory and antioxidant activity of Sri Lankan medicinal plants for novel cosmeceuticals. *Industrial Crops and Products*, 111(October 2017), 597–605. <https://doi.org/10.1016/j.indcrop.2017.11.019>
- Lotulung, P. D. N., Mozef, T., Risdian, C., & Darmawan, A. (2014). In vitro antidiabetic activities of extract and isolated flavonoid compounds from Artocarpus altilis (Parkinson) Fosberg. *Indonesian Journal of Chemistry*, 14(1), 7–11. <https://doi.org/10.22146/ijc.21261>
- Mahadi Abdul Jamil, M., Ganeson, S., Buhari Mammam, H., & Abdul Wahab, R. (2018). Artocarpus altilis extract effect on cervical cancer cells. *Materials Today: Proceedings*, 5(7), 15559–15566. <https://doi.org/10.1016/j.matpr.2018.04.163>

- Mai, N. T. T., Hai, N. X., Phu, D. H., Trong, P. N. H., & Nhan, N. T. (2012). Three new geranyl aurones from the leaves of *Artocarpus altilis*. *Phytochemistry Letters*, 5(3), 647–650. <https://doi.org/10.1016/j.phytol.2012.06.014>
- Mazimba, O., Majinda, R. R. T., & Motlhanka, D. (2011). Antioxidant and antibacterial constituents from *Morus nigra*. *African Journal of Pharmacy and Pharmacology*, 5(6), 751–754. <https://doi.org/10.5897/AJPP11.260>
- Mozef, T., Risdian, C., Sukandar, E. Y., & Soemardji, A. A. (2015). Bioactivity of Ethyl Acetate Fraction from the Leaves of “Sukun” (*Artocarpus Altilis* (Parkinson) Fosberg) in Preventing Atherosclerosis. *Procedia Chemistry*, 16, 106–112. <https://doi.org/10.1016/j.proche.2015.12.037>
- Namdaung, U., Athipornchai, A., Khammee, T., Kuno, M., & Suksamrarn, S. (2017). 2-Arylbenzofurans from *Artocarpus lakoocha* and methyl ether analogs with potent cholinesterase inhibitory activity. *European Journal of Medicinal Chemistry*. <https://doi.org/10.1016/j.ejmech.2017.10.019>
- Priska, M., Peni, N., & Carvallo, L. (2019). Phytochemicals Screening and Antioxidant Effectiveness of Garlic (*Allium sativum*) from Timor Island. *Biosaintifika: Journal of Biology & Biology Education*, 11(1), 1–7. <https://doi.org/10.15294/biosaintifika.v11i1.17313>
- Septama, A. W., & Panichayupakaranant, P. (2015). Antibacterial assay-guided isolation of active compounds from *Artocarpus heterophyllus* heartwoods. *Pharmaceutical Biology*, 53(11), 1608–1613. <https://doi.org/10.3109/13880209.2014.996819>
- Shamaun, S. S., Rahmani, M., Hashim, N. M., Ismail, H. B. M., Sukari, M. A., Lian, G. E. C., & Go, R. (2010). Prenylated flavones from *Artocarpus altilis*. *Journal of Natural Medicines*, 64(4), 478–481. <https://doi.org/10.1007/s11418-010-0427-4>
- Sikarwar, M. S., Hui, B. J., Subramaniam, K., Valeisamy, B. D., Yean, L. K., & Balaji, K. (2014). A review on *Artocarpus altilis* (Parkinson) Fosberg (breadfruit). *Journal of Applied Pharmaceutical Science*, 4(8), 91–97. <https://doi.org/10.7324/JAPS.2014.40818>
- Somashekhar, M., Nayeem, N., & Sonnad, B. (2013). A review on family Moraceae (Mulberry) with a focus on *Artocarpus* species. *World J. Pharm. Pharm. Sci.*
- Swinney, D. C. (2011). Molecular Mechanism of Action (MMoA) in Drug Discovery. In *Annual Reports in Medicinal Chemistry* (1st ed., Vol. 46). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-386009-5.00009-6>
- Thuy, T. T. (2016). In vivo anticancer activity of maesopsin 4-O- β -glucoside isolated from leaves of *Artocarpus tonkinensis* A. Chev. Ex Gagnep. *Asian Pacific Journal of Tropical Medicine*. <https://doi.org/10.1016/j.apjtm.2016.03.012>
- Tsimogiannis, D., Samiotaki, M., Panayotou, G., & Oreopoulou, V. (2007). Characterization of Flavonoid Subgroups and Hydroxy Substitution by HPLC-MS/MS. *Molecules*, 593–606.
- Wang, Y., Deng, T., Lin, L., Pan, Y., & Zheng, X. (2008). Bioassay-guided Isolation of Antiatherosclerotic Phytochemicals from *Artocarpus altilis*. *Phytotherapy Research*, 22(4), 544–549. <https://doi.org/10.1002/ptr>
- Wei, B. L., Weng, J. R., Chiu, P. H., Hung, C. F., Wang, J. P., & Lin, C. N. (2005). Antiinflammatory flavonoids from *Artocarpus heterophyllus* and *Artocarpus*

- communis. *Journal of Agricultural and Food Chemistry*, 53(10), 3867–3871.
<https://doi.org/10.1021/jf047873n>
- Weng, J., Chan, S., Lu, Y., Lin, H., Ko, H., & Lin, C. (2006). *Antiplatelet prenylflavonoids from Artocarpus communis.* 67, 824–829.
<https://doi.org/10.1016/j.phytochem.2006.01.030>
- Yuan, W., Yuan, J., Peng, J., Ding, Y., Zhu, J., & Ren, G. (2017). Flavonoids from the roots of Artocarpus heterophyllus. *Fitoterapia*, 117, 133–137.
<https://doi.org/10.1016/j.fitote.2017.01.016>

