

DAFTAR PUSTAKA

- Achmad, S. A., Hakim, E. H., Juliawaty, L. D., Makmur, L., Sutoyo, S., Aimi, N., & Ghisalberti, E. L. (1996). A new prenylated flavone from *Artocarpus champeden*. *Journal of Natural Products*, 59(9), 878–879. <https://doi.org/10.1021/np960330q>
- Arung, E. T., Wicaksono, B. D., Handoko, Y. A., Kusuma, I. W., Yulia, D., & Sandra, F. (2009). Anti-cancer properties of diethylether extract of wood from Sukun (*Artocarpus altilis*) in human breast cancer (T47D) cells. *Tropical Journal of Pharmaceutical Research*, 8(4), 317–324. <https://doi.org/10.4314/tjpr.v8i4.45223>
- Asih, I. A. R. A., Sudiarta, I. W., & Suci, A. A. W. (2015). Aktivitas Antioksidan Senyawa Golongan Flavonoid Ekstrak Etanol Daging Buah Terong Belanda (*Solanum betaceum* Cav.). *Jurnal Kimia*, 9(1), 35–40.
- Badarinath, A. V., Rao, K. M., Chetty, C. M. S., Ramkanth, S., Rajan, T. V. S., & Gnanaprakash, K. (2010). A Review on In-vitro Antioxidant Methods: Comparisons, Correlations and Considerations, 2(2), 1276–1285.
- Boonlaksiri, C., Oonanant, W., Kongsaree, P., Kittakoop, P., Tanticharoen, M., & Thebtaranonth, Y. (2000). An antimalarial stilbene from *Artocarpus integer*. *Phytochemistry*, 54(4), 415–417. [https://doi.org/10.1016/S0031-9422\(00\)00074-1](https://doi.org/10.1016/S0031-9422(00)00074-1)
- de Almeida Lopes, M. M., de Souza, K. O., & de Oliveira Silva, E. (2018). Cempedak— *Artocarpus champeden*. *Exotic Fruits*, 121–127. <https://doi.org/10.1016/B978-0-12-803138-4.00017-4>
- Ersam, T., Achmad, S. A., Ghisalberti, E. L., Hakim, E. H., Makmur, L., & Syah, Y. M. (2002). A new isoprenylated chalcone, artoindonesianin J, from the root and tree bark of *Artocarpus bracteata*. *Journal of Chemical Research-S*, (4), 186–187.
- Fessenden, R. ., & Fessenden, J. . (1986). *Kimia Organik*. Jakarta: Erlangga.
- Garcia, E. J., Cadorin Oldoni, T. L., de Alencar, S. M., Reis, A., Loguercio, A. D., & Miranda Grande, R. H. (2012). Antioxidant activity by DPPH assay of potential solutions to be applied on bleached teeth. *Brazilian Dental Journal*, 23(1), 22–27. <https://doi.org/10.1590/S0103-64402012000100004>
- Grotewold, E. (2006). *The science of flavonoids. The Science of Flavonoids*. <https://doi.org/10.1007/978-0-387-28822-2>
- Hakim, E. H., Achmad, S. A., Juliawaty, L. D., Makmur, L., Syah, Y. M., Aimi, N., ... Ghisalberti, E. L. (2007). Prenylated flavonoids and related compounds of the Indonesian *Artocarpus* (Moraceae) (*Journal of Natural Medicines* 60, (161-184) DOI: 10.1007/s11418-006- 0048-0). *Journal of Natural Medicines*, 61(2), 229. <https://doi.org/10.1007/s11418-006-0106-7>

- Hakim, E. H., Mar, E., Mujahidin, D., Achmad, S. A., Ghisalberti, E. L., & Makmur, L. (1998). Artokarpin dan Heteroflavon-A, dua senyawa flavonoid bioaktif dari *Artocarpus champeden*, 0–5.
- Hanh, N. H., & Hoa, L. T. P. (2015). Tyrosinase inhibitors from the heartwood of Vietnamese *Artocarpus heterophyllus*, 5(4), 1–4.
- Heyney, K. (1987). *Tumbuhan Berguna Indonesia* (II). Jakarta: Yayasan Sarana Wana Jaya.
- Jayasinghe, L., Rupasinghe, G. K., Hara, N., & Fujimoto, Y. (2006). Geranylated phenolic constituents from the fruits of *Artocarpus nobilis*. *Phytochemistry*, 67(13), 1353–1358. <https://doi.org/10.1016/j.phytochem.2006.04.011>
- Khan, M. R., Omoloso, A. D., & Kihara, M. (2003). Antibacterial activity of *Artocarpus heterophyllus*. *Fitoterapia*, 74(5), 501–505. [https://doi.org/10.1016/S0367-326X\(03\)00120-5](https://doi.org/10.1016/S0367-326X(03)00120-5)
- Kochummen, K. (1987). *Moracea in tree flora of Malaya* (II). Kepong: Forest Research Institute.
- Kristanti, A., Aminah, N., Tanjung, M., & Kurniadi, B. (2008). *Buku Ajar Fitokimia*. Surabaya: Universitas Airlangga.
- Mabruroh, E. Q., Mursiti, S., & Kusumo, E. (2019). Indonesian Journal of Chemical Science Isolasi dan Identifikasi Senyawa Flavonoid dari Daun Murbei (*Morus alba* Linn), 8(1).
- Margianasari, A. F., Kusumahastuti, S. W., Junaedi., Parimin., Dahlan, D., & Sodikin. (2009). *Nangkadak Mekarsari*. (Junaedi, F. . Kerong, & Nasrul, Eds.). Bogor: Taman Wisata Mekarsari.
- Molyneux, P. (2004). The Use of the Stable Free Radical Diphenylpicryl-hydrazyl (DPPH) for Estimating Antioxidant Activity. *Songklanakarin Journal of Science and Technology*, 26(December 2003), 211–219. <https://doi.org/10.1287/isre.6.2.144>
- Nomura, T., Hano, Y., & Aida, M. (1998). Isoprenoid-substituted flavonoids from *Artocarpus* plants (moraceae). *Heterocycles*, 47(2), 1179–1205. [https://doi.org/10.3987/REV-97-SR\(N\)9](https://doi.org/10.3987/REV-97-SR(N)9)
- Seibert, B., & P.C.M, J. (1997). *Buah-buahan yang dapat dimakan. Sumber Daya Nabati Asia Tenggara (PROSEA)* (II). Jakarta: Gramedia.
- Somashekhar., M., Nayeem., N., & B, S. (2013). A review on family Moraceae (Mulberry) with a focus on *Artocarpus* species. *World Journal Of Pharmacy And Pharmaceutical Sciences*, 2, 2614–2621.
- Syamsuhidayat, S. S., & Hutapea, J. R. (1991). *Inventaris Tanaman Obat Indonesia* (II). Jakarta: Departemen Kesehatan RI, Badan Penelitian dan Pengembangan Kesehatan.

- Weng, J. R., Chan, S. C., Lu, Y. H., Lin, H. C., Ko, H. H., & Lin, C. N. (2006). Antiplatelet prenylflavonoids from *Artocarpus communis*. *Phytochemistry*, 67(8), 824–829. <https://doi.org/10.1016/j.phytochem.2006.01.030>
- Widyawaruyanti, A., Subehan, Kalauni, S. K., Awale, S., Nindatu, M., Zaini, N. C., ... Kadota, S. (2007). New prenylated flavones from *Artocarpus champeden*, and their antimalarial activity in vitro. *Journal of Natural Medicines*, 61(4), 410–413. <https://doi.org/10.1007/s11418-007-0153-8>
- Zschocke, S., & Van Staden, J. (2000). Cryptocarya species - Substitute plants for *Ocotea bullata*. A pharmacological investigation in terms of cyclooxygenase-1 and -2 inhibition. *Journal of Ethnopharmacology*, 71(3), 473–478. [https://doi.org/10.1016/S0378-8741\(00\)00183-5](https://doi.org/10.1016/S0378-8741(00)00183-5)

